

AD _____

Grant Number DAMD17-97-1-7009

TITLE: Emergency Interventions After Severe Traumatic Brain
Injury in Rats: Effect on Neuropathology and Functional Outcome

PRINCIPAL INVESTIGATOR: Patrick M. Kochanek, M.D.

CONTRACTING ORGANIZATION: University of Pittsburgh
Pittsburgh, Pennsylvania 15260

REPORT DATE: January 1998

TYPE OF REPORT: Annual

DTIC QUALITY INSPECTED 3

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19980219 151

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 1998	3. REPORT TYPE AND DATES COVERED Annual (15 Dec 96 - 14 Dec 97)	
4. TITLE AND SUBTITLE Emergency Interventions After Severe Traumatic Brain Injury in Rats: Effect on Neuropathology and Functional Outcome			5. FUNDING NUMBERS DAMD17-97-1-7009	
6. AUTHOR(S) Patrick M. Kochanek, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Pittsburgh Pittsburgh, Pennsylvania 15260			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200) Traumatic brain injury (TBI) contributes to combat casualty morbidity and mortality. Our hypothesis is that optimal manipulation of practical interventions applicable in the emergency treatment of severe TBI (respiratory management, temperature control, and sedation) can reduce brain injury in a rat model of brain contusion, and thereby improve functional and neuropathological outcome. In the first year of funding, we addressed the first Technical Objective of our proposal - to perform a comprehensive study of the effects of mechanical ventilation strategies on outcome. We found that aggressive hyperventilation applied for 4 h immediately after injury is detrimental (vs normal ventilation), and increases neuronal death in vulnerable brain regions. Also, to set the stage for the evaluation of therapies targeting improved outcome after TBI (proposed in Technical Objectives 2-4), the severity of the insult was increased in our model. This was done by more accurately simulating the field scenario (adding a secondary insult). Finally, another injury station was established and a technician was trained to perform the studies proposed in y 2-3. Dr. Michael Forbes, a fellow completed his training during this year and was the first author of the manuscript described above.				
14. SUBJECT TERMS Head Injury, Hyperventilation, Hypothermia, Trauma, Apoptosis, Excitotoxicity, Sedation, Oxygen			15. NUMBER OF PAGES 48	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

✓
Where copyrighted material is quoted, permission has been obtained to use such material.

✓
Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

✓
Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


✓
In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

 Jan 12, 1988

PI - Signature Date

Table of Contents

SF 298	2
FOREWORD	3
TABLE OF CONTENTS	4
INTRODUCTION	5
BODY	5
CONCLUSIONS	13
REFERENCES	14
APPENDICES	15

(5) INTRODUCTION

In our application, we highlighted the fact that traumatic brain injury (TBI) is an important contributor to combat casualty morbidity and mortality. We also stated that although progress has been made in the development of rodent models of TBI (such as the controlled cortical impact (CCI) model), most of the studies with these models have focused on molecular mechanisms of damage. Because there has been a paucity of application of **practical emergency interventions** in TBI models, we felt that it was essential to address this deficiency and that this strategy could have important implications for field and emergency management of both soldiers and civilians with severe TBI. Our overall **hypothesis** is that optimal manipulation of practical interventions applicable in the emergency treatment of severe TBI (respiratory management, temperature control, and sedation) can reduce secondary brain injury in a rat model of brain contusion, and thereby improve functional and/or neuropathological outcome.

In the first year of funding, we addressed the most important aspect of the **first Technical Objective** of our proposal -namely- to perform a comprehensive study of the effects of mechanical ventilation strategies (as applied by the first responder in the field) on both functional and neuropathological outcome in our model. **We found that aggressive, prophylactic hyperventilation (HV) applied for 4 h immediately after injury is detrimental (vs ventilation to a normal PaCO₂, normal ventilation [NV]), and leads to an increase in the amount of neuronal death in selectively vulnerable brain regions.** This study, is now *in press* as a full manuscript in the *Journal of Neurosurgery* (1), (also see Appendix #1). We were pleased that the reviewers indicated that this was an important study which would be cited often.

In addition, to set the stage for the evaluation of therapies targeting improvement in outcome after severe TBI (as proposed studies in technical objectives 2-4), it appeared that it would be optimal to increase the severity of the insult in our model. This was done by attempting to more accurately simulate the field scenario – i.e., adding a 30 min period of moderate hypoxemia to the insult. The characterization of that model for our future studies will also be described below.

During the first year of funding, Henry Alexander, an experienced technician assumed the technical duties of the injury model and has successfully learned the model to perform all of the subsequent injury studies in years 2 and 3. Also, a new injury device and station were purchased and is in operation for these studies. Finally, Dr. Michael Forbes, a fellow in Pediatric Critical Care Medicine completed his training during this first year of funding and was the team leader on our study assessing the effect of HV in our model. He was the first author of the manuscript describing that work. Dr. Forbes is now Associate Director of the Pediatric Intensive Care Unit at Allegheny General Hospital in Pittsburgh.

(6) BODY

(a) Technical Objective 1: Mechanical ventilation strategies after severe TBI in rats

For over two decades, HV has been one of the most utilized strategies in the management of TBI. Laboratory and clinical studies, however, have verified that early after TBI, there is usually a state of reduced cerebral perfusion that may increase vulnerability to secondary injury. HV reduces intracranial hypertension by reducing cerebral blood volume; however, this generally is accompanied by a reduction in cerebral blood flow. A recent clinical study

suggested that HV may worsen outcome after TBI. However, in the field or during the initial stabilization, HV is often used (either planned or iatrogenically) and the first blood gas of patients in the emergency room can reveal significant hypocarbia. Using the CCI model in rats, we tested the effect of 4 h of aggressive HV (vs NV), beginning immediately after injury, on functional and neuropathological outcome.

Methods:

All of the protocols for the studies outlined below were approved by the Animal Care and Use Committee of the University of Pittsburgh. Intubated, mechanically ventilated, isoflurane-anesthetized male Sprague-Dawley rats ($n = 26$) were subjected to CCI (4 m/s, 2.5 mm depth of deformation) and randomized after 10 min to either HV [$n = 13$, $P_aCO_2 = 20.3 \pm 0.7$ mm Hg] or NV [$n = 13$, $P_aCO_2 = 34.9 \pm 0.3$ mm Hg] for 5 h. Beam balance and Morris water maze (MWM) performance latencies were measured in eight rats from each group on d 1-5 and 7-11 post CCI, respectively. Rats were killed at 14 d. Serial coronal sections of brain were studied for contusion volume and hippocampal neuron counting [CA1, CA3] by a blinded observer.

Results:

HV was readily achieved and could be sustained for 4 h in our model, and produced the anticipated systemic alkalosis (Figure 1). In addition, other variables could be tightly controlled for 4 h in our model (Figure 1).




Figure 1. Entire time course of (A) P_aCO_2 (mm Hg), (B) arterial pH, (C) Mean arterial blood pressure (MABP, mm Hg), and (D) brain temperature ($^{\circ}C$) in all rats treated with either NV (\blacktriangle , $n = 13$) or HV (\blacksquare , $n = 13$) after CCI. * $p < 0.05$ for NV vs HV. Data are mean \pm SEM.

Mortality rates were similar in both groups (2/13 vs 3/13, NV vs HV, respectively, *NS*). There were no differences between groups in mean arterial blood pressure, brain temperature, and serum glucose concentration. There were no differences between groups in either performance latencies for both beam balance (Figure 2) and MWM (Figure 3) or contusion volume (Figure 4).




Figure 2. Mean beam balance performance latencies (mean \pm SEM, in sec) in rats before and on d 1-5 after CCI (4 m/s, 2.5 mm cortical deformation depth). Analysis of variance with repeated measures revealed no difference in duration of balance maintained between the two groups. (\blacktriangle , NV, $n = 8$; \blacksquare , HV, $n = 8$). Data are mean \pm SEM.




Figure 3. MWM performance latency to find a hidden platform (mean \pm SEM, in sec) by rats on d 7-11 after CCI. There was no between group difference (\blacktriangle , NV, $n = 8$; \blacksquare , HV, $n = 8$) when performances were compared using ANOVA with repeated measures. Data are mean \pm SEM.




Figure 4. Graph depicting mean lesion area (left y-axis, mm²) vs distance from occiput (mm) measured 14 d after CCI (NV, open bars, n = 11, HV, closed bars, n = 10). Contusion volume (mm³) was calculated as the sum of these areas in each group and is depicted as cumulative volume (right y-axis) in the NV, ▲, and HV, ■, groups. There was no difference between groups in contusion volume (27.8 ± 5.1 vs 27.8 ± 3.1 mm³ NV vs HV, mean \pm SEM).

However, in brain sections through the center of the contusion, hippocampal neuronal survival in HV reduced the number of surviving hippocampal CA3 neurons (29.7 [24.2 - 31.7] vs 19.9 [17.0 - 23.7] cells/high power field (NV vs HV, median [25th-75th percentiles] * $p < 0.05$, Mann-Whitney Rank Sum Test, Figure 5). In contrast to the detrimental effect on CA3 neurons, CA1 neuronal death was not increased by aggressive HV.




Figure 5. Box plots representing the number of surviving CA1 and CA3 hippocampal neurons in coronal brain sections through the center of the lesion in the hemisphere ipsilateral to the contusion. Cells were counted 14 d after injury. The median line is placed within the shaded 25th - 75th range. There was a reduction in the number of surviving CA3 hippocampal neurons after injury comparing NV and HV groups (29.7 [24.2 - 31.7] vs 19.9 [17.0 - 23.7], cells/high power field (hpf), * $p < 0.05$, Mann-Whitney rank sum test).

Conclusion:

Aggressive HV early after TBI augments CA3 hippocampal neuronal death; however, it did not impair functional outcome or expand the contusion. These data suggest that CA3 hippocampal neurons are selectively vulnerable to the effects of HV early after TBI. The injudicious application of HV early after TBI, such as in the field, may contribute to secondary neuronal injury.

Comment:

Hippocampal CA3 neurons are selectively vulnerable to delayed neuronal death after TBI. The mechanisms underlying this process remain speculative. Potential mechanisms include ischemia, TBI-induced excitotoxicity, apoptosis, and inflammation. We previously demonstrated that the hippocampus and cortex ipsilateral to the impact have marked flow reduction (at least 60%) at 2 h after TBI in the CCI model (2). CBF approaches ischemic levels in the core of the contusion at 2 h after injury. Although we have not evaluated the status of reactivity of the cerebral circulation to changes in PaCO₂ at 2 h after TBI in this model, we have reported that CO₂ reactivity is impaired, although still present (62-71% of baseline) in and around the contusion at 24 h after CCI in rats (3).

HV produces cerebral vasoconstriction and alkalosis. Alkalosis exacerbates N-methyl-D-aspartate (NMDA)-receptor mediated neurotoxicity. As a result of aggressive HV, the rats in our study were quite alkalotic as indicated by arterial pH measurements. Alkalosis appears to have deleterious effects on neurons. It could also be that the combined effect of alkalosis and further flow reduction by HV is deleterious in regions vulnerable to excitotoxicity such as CA3. Early, aggressive or prophylactic HV, therefore, in the context of reduced CBF, may exacerbate excitotoxic mechanisms and augment neuronal death.

Aggressive HV in the early low flow period did not worsen functional outcome or expand the contusion. The cognitive deficits in this model are modest. Indeed, to test therapies targeting an improvement in outcome, we may need a more severe injury (see below). Additional unilateral or bilateral hippocampal damage may be necessary to create more marked functional deficits. CA3 damage alone may not mediate post-TBI MWM deficits. However, hippocampal damage and memory deficits are common after TBI in humans, and exacerbation of neuronal death in any brain region would be highly undesirable.

This study does not completely address the uncommon situation where early after severe head injury marked intracranial hypertension is observed. HV may, in fact, be life saving in the setting of impending herniation. Similarly, we did not measure ICP or titrate ventilation to control cerebral perfusion pressure, and we evaluated only one level of HV and injury severity. We did not attempt to model the clinical scenario of optimal titration of ventilation when ICP is increased. Rather, we chose to evaluate the field setting and apply the worst case scenario, aggressive HV during the early post-trauma period when flow is already low and excitotoxicity is peaking. Our study does, however, show that HV is associated with a tangible risk to vulnerable neurons. To our knowledge, this is the first *in vivo* study demonstrating that HV can augment neuronal injury after TBI. This suggests that there is indeed a trade-off associated with this intervention.

Recommendation

We have shown that aggressive, early HV after TBI augments neuronal death in CA3 hippocampus in a rat model of cerebral contusion. The further reduction of CBF with HV during the low cerebral blood flow state immediately after severe TBI coupled with alkalosis may increase the vulnerability of selected neurons to damage. The results of this study reinforce the meticulous attention necessary to prevent secondary injury after TBI. A risk in the use of HV is demonstrated. Unless signs of impending herniation (unilateral pupillary dilation, hypertension, bradycardia) are present, this study supports the targeting of normocarbia for mechanical ventilation in the emergency stabilization of the brain trauma victim.

(b) A field scenario of severe TBI for the evaluation of therapies proposed in Technical Objectives 2-4.

Using our standard CCI model, neuronal death in selectively vulnerable regions and MWM deficits are present but modest. In tat model, we able to nicely demonstrate exacerbation of damage with a deleterious strategy, namely HV. However, in technical objectives 2-4, our goal is to define strategies (hypothermia, anesthetics, anti-excitotoxic therapies) that will mitigate damage. Thus, the severity of damage must be increased, both from the standpoint of both hippocampal neuronal death and MWM deficit, to achieve this goal. Recently, in studies separate from this application, we published a variation of our CCI model that was designed to

increase the amount of hippocampal damage without totally destroying the hippocampus (and making it impossible to resuscitate)(4). This was achieved by adding a 30 min period of moderate hypoxemia ($FiO_2 = 0.11$) which also results in accompanying mild hypotension. This mimics the secondary insults in head injury victims so commonly seen in the field. In addition, in studies by our group separate from this application (5), we reported that both necrotic and apoptotic neuronal death is seen in this new variant of the CCI model. To be certain that this model would be suitable for technical objectives 2-4, it was essential to determine if the insult was accompanied by a significant MWM deficit.

Methods:

All of the protocols for the studies outlined below were approved by the Animal Care and Use Committee of the University of Pittsburgh. Intubated, mechanically ventilated, isoflurane-anesthetized male Sprague-Dawley rats ($n = 20$) were subjected to CCI (4 m/s, 2.5 mm depth of deformation) to the left parietal cortex using either a vertical or angled impact. Immediately after injury, the FiO_2 was reduced to 0.11 (inhalational anesthesia maintained constant by the addition of N_2 to the ventilator circuit). At 5 min after reducing the FiO_2 and at the completion of the 30 min secondary hypoxemic insult, a blood gas is obtained to document the level of hypoxemia achieved, and then the FiO_2 is increased. Shams were subjected to all surgical procedures, but no insult (i.e., neither CCI nor hypoxemia). After the recovery periods, catheters were removed and anesthesia was discontinued. Rats were weaned from mechanical ventilation, extubated, and returned to their cages until further study. Motor and cognitive outcome were assessed as previously described.

Results:

Both vertical and angled impacts resulted in significant motor and cognitive deficits as assessed by beam balance, and MWM paradigms (Figures 6, and 7, respectively).




Figure 6. Mean beam balance performance latencies (mean \pm SEM, in sec) in rats before and on d 1-5 after either vertical or angled CCI with secondary hypoxemic insult. Analysis of variance with repeated measures revealed no difference in duration of balance maintained between the two insults. Both insults were significantly different from sham.




Figure 7. MWM performance latency to find a hidden platform (mean \pm SEM, in sec) by rats on d 14-21 after either vertical or angled CCI with secondary hypoxemic insult. Analysis of variance with repeated measures revealed no difference between the two insults. Both insults were significantly different from sham.

Conclusion:

Combined with our prior publications showing both a well defined contusion and neuronal death by both apoptosis and necrosis in this model (4,5), the functional deficits produced with either a vertical or angled insult set the stage for studies proposed in Technical Objectives 2-4. We have chosen to use the vertical insult with hypoxemia, since all of our initial studies with HV used a vertical impact. These will be addressed in years 2-3 of the funding period. In accordance with this plan, we are currently evaluating the effect of hypothermia in this model of CCI with a secondary insult (as outlined in Technical Objective #2). We plan to address Technical Objective 2 and part of Technical Objective 3 in funding year 2.

(7) CONCLUSION

In our work during the first year of funding addressing Technical Objective #1, we demonstrated that aggressive HV early after TBI augments CA3 hippocampal neuronal death. These data suggest that CA3 hippocampal neurons are selectively vulnerable to the effects of HV early after TBI. The injudicious application of HV early after TBI, such as in the field, may contribute to secondary neuronal injury. As previously discussed, the results of this study reinforce the meticulous attention necessary to prevent secondary injury after TBI. A risk in the use of HV is demonstrated. Unless signs of impending herniation (unilateral pupillary dilation, hypertension, bradycardia) are present, this study supports the targeting of normocarbica for mechanical ventilation in the emergency stabilization of the brain trauma victim. Finally, by adding a secondary insult to our injury model, we have set the stage to address the optimal

application of treatments to improve outcome as outlined in Technical Objectives 2-4, and those studies are underway.

(8) REFERENCES

1. Forbes ML, Clark RSB, Dixon CE, Graham SH, Marion DW, DeKosky ST, Schiding JK, Kochanek PM: Hyperventilation Early After Controlled Cortical Impact Augments Neuronal Death in CA3 Hippocampus. *J Neurosurg* (in press).
2. Kochanek PM, Marion D, Zhang W, Schiding J, White M, Palmer A, Clark RSB, O'Malley M, Styren S, Ho C, DeKosky S: Severe Cortical Impact in Rats: Assessment of Cerebral Edema, Blood Flow, and Contusion Volume. *J Neurotrauma* 12:1015-1025, 1995.
3. Forbes ML, Hendrich KS, Kochanek PM, Williams DS, Schiding JK, Wisniewski SR, Kelsey SF, DeKosky ST, Graham SH, Marion DW, Ho C: Assessment of Cerebral Blood Flow and CO₂ Reactivity After Controlled Cortical Impact By Perfusion Magnetic Resonance Imaging Using Arterial Spin Labeling in Rats. *J Cereb Blood Flow Metab* 17:865-874, 1997.
4. Clark RSB, Kochanek PM, Dixon CE, Chen M, Marion DW, Heineman S, DeKosky ST, Graham SH: Early Neuropathologic Effects of Mild or Moderate Hypoxemia After Controlled Cortical Impact Injury in Rats. *J Neurotrauma* 14:179-189, 1997.
5. Clark RSB, Chen J, Watkins SC, Kochanek PM, Chen M, Stetler RA, Graham SH: Apoptosis-Suppressor Gene *bcl-2* Expression After Traumatic Brain Injury in Rats. *J Neurosci* 17:9172-9182, 1997.

APPENDIX

Journal of Neurosurgery

1224 WEST MAIN STREET, SUITE 450
CHARLOTTESVILLE, VIRGINIA 22903

TELEPHONE: (804) 924-5503 • FAX: (804) 924-2702 • E-MAIL: jneuro@virginia.edu

Editor

John A. Jane, M.D., Ph.D.

Managing Editor

Keller Kaufman-Fox

Editorial Staff

Arthur W. Chapin
Melba C. Christy
Jo Ann M. Eliason
Donna M. Goodrich
Evelyn C. Kessler
Cathleen A. Nielan
Brenda H. Shifflett
Margie A. Shreve
Charlotte D. Tolar
Dorothy L. Turner
Patricia Whitton

Associate Editor for

Neurosurgical Focus
Martin H. Weiss, M.D.
Los Angeles, California

Editorial Board

Co-Chairman
Howard M. Eisenberg, M.D.
Baltimore, Maryland

Co-Chairman
Julian T. Hoff, M.D.
Ann Arbor, Michigan

John P. Girvin, M.D., Ph.D.
London, Ontario, Canada

Donald P. Becker, M.D.
Los Angeles, California

Edward R. Laws, Jr., M.D.
Charlottesville, Virginia

Robert A. Ratcheson, M.D.
Cleveland, Ohio

M. Peter Heilbrun, M.D.
Salt Lake City, Utah

Charles J. Hodge, Jr., M.D.
Syracuse, New York

Edward H. Oldfield, M.D.
Bethesda, Maryland

Ralph G. Dacey, Jr., M.D.
St. Louis, Missouri

H. Richard Winn, M.D.
Seattle, Washington

Lawrence F. Marshall, M.D.
San Diego, California

Edward C. Benzel, M.D.
Albuquerque, New Mexico

October 23 1997

Patrick M. Koshanek, M.D.
Safer Center for Resuscitation Research
3434 Fifth Avenue
Pittsburgh, PA 15260

RE: Manuscript #8160

Dear Doctor Koshanek:


Thank you for your responsive revision of the paper entitled *Hyperventilation early after controlled cortical impact augments neuronal death in CA3 hippocampus*. It is now accepted for publication in the *Journal of Neurosurgery*.

Please send us an original revised title page that includes full names of all co-authors (first, middle and last) at your earliest convenience. We will then proceed with publication. We look forward to receiving the completed copyright form.

Our editorial staff will be in contact with you as your manuscript is prepared for publication. We would appreciate your timely response to queries that we have, as this plays an important part in determining to which issue your paper is assigned.

This is an important and excellent study. Thank you for sending it to us.

Sincerely,



John A. Jane, M.D., Ph.D.
Editor

Journal of Neurosurgery

JAJ:mas

Hyperventilation Early After Controlled Cortical Impact Augments Neuronal Death in CA3 Hippocampus

Forbes ML, MD¹, Clark RSB, MD^{1,4}, Dixon CE, PhD^{1,6}, Graham SH, MD, PhD⁵,
Marion DW, MD^{2,6}, DeKosky ST, MD^{2,7}, Schiding JK, BS¹,
Kochanek PM, MD^{1,2,3,4}

¹*Safar Center for Resuscitation Research*, ²*The University of Pittsburgh Brain
Trauma Research Center and Departments of* ³*Anesthesiology/Critical Care
Medicine*, ⁴*Pediatrics*, ⁵*Neurology*, ⁶*Neurosurgery* and ⁷*Psychiatry*,
University of Pittsburgh, Pgh, PA 15260

Address correspondence and reprint requests to:

Patrick M. Kochanek, M.D.
Safar Center for Resuscitation Research
3434 Fifth Avenue
Pittsburgh, PA 15260
Telephone: (412) 383-1900
Fax: (412) 624-0943
Internet: kochanek@smtp.anes.upmc.edu

Sources of support:

US Army Medical Research Acquisition Activity #DAMDI17-97-1-7009 (PK)
The Laerdal Foundation (PK)
The National Institutes of Health: NS30318 (DM), NS3198 (CED)
Children's Hospital of Pittsburgh Seed Grant (MF)
Department of Veterans Medical Affairs Merit Review Program (SG)

Running Head: Hyperventilation after TBI augments neuronal death

Key Words: Head injury, rat, hyperventilation, alkalosis, hippocampus

Manuscript: #8160

Abstract

Introduction: Minimizing secondary injury after severe traumatic brain injury (TBI) is the primary goal of cerebral resuscitation. For over two decades, hyperventilation (HV) has been one of the most utilized strategies in the management of TBI. Laboratory and clinical studies, however, have verified a post-TBI state of reduced cerebral perfusion that may increase vulnerability to secondary injury. In addition, a clinical study suggested that HV may worsen outcome after TBI. Using the controlled cortical impact (CCI) model in rats, we tested the hypothesis that aggressive HV applied immediately after TBI would worsen functional outcome, expand the contusion, and promote neuronal death in selectively vulnerable hippocampal neurons.

Methods: Intubated, mechanically ventilated, isoflurane-anesthetized male Sprague-Dawley rats ($n = 26$) were subjected to CCI (4 m/s, 2.5 mm depth of deformation) and randomized after 10 min to either HV [$n = 13$, $P_aCO_2 = 20.3 \pm 0.7$ mm Hg] or normal ventilation [NV; $n = 13$, $P_aCO_2 = 34.9 \pm 0.3$ mm Hg] for 5 h. Beam balance and Morris water maze (MWM) performance latencies were measured in eight rats from each group on d 1-5 and 7-11 post CCI, respectively. Rats were killed at 14 d. Serial coronal sections of brain were studied for contusion volume and hippocampal neuron counting [CA1, CA3] by a blinded observer.

Results: Mortality rates were similar in both groups (2/13 vs 3/13, NV vs HV, respectively, *NS*). There were no differences between groups in mean arterial blood pressure, brain temperature, and serum glucose concentration. There were no differences between groups in either performance latencies for both beam balance and MWM or contusion volume (27.8 ± 5.1 mm³ vs 27.8 ± 3.3 mm³; NV vs HV, respectively, *NS*). In brain sections through the center of the contusion, hippocampal neuronal survival in the CA1 region was similar in both groups; however, HV reduced the number of surviving hippocampal CA3 neurons (29.7 [24.2

- 31.7] vs 19.9 [17.0 - 23.7] cells/high power field, NV vs HV, median [25th-75th percentiles] $*p < 0.05$, Mann-Whitney Rank Sum Test).

Conclusion: Aggressive HV early after TBI augments CA3 hippocampal neuronal death; however, it did not impair functional outcome or expand the contusion. These data suggest that CA3 hippocampal neurons are selectively vulnerable to the effects of HV after TBI. Further studies delineating the mechanisms underlying these effects are needed. The injudicious application of HV early after TBI may contribute to secondary neuronal injury.

Introduction

Traumatic brain injury (TBI) is often complicated by malignant intracranial hypertension,³¹ which is associated with high mortality and has been managed using a combination of therapies including osmotherapy, diuretics, sedation, neuromuscular blockade, optimization of cerebral perfusion pressure, and hyperventilation (HV).^{5,8,31,38,50} HV has been an integral part of the clinical armamentarium in the management of severe TBI for over 20 years.¹¹ HV rapidly reduces cerebral blood flow (CBF) and cerebral blood volume (CBV) in areas of brain with intact CO₂ autoregulation, providing one therapeutic option in the management of TBI complicated by malignant intracranial hypertension.^{1,33,41}

Recent studies, however, have defined a state of reduced CBF early after TBI in humans^{2,30} and animals^{4,19,24,45,56,57}, particularly in the first 8 h after TBI. Some have hypothesized that the brain is more vulnerable to secondary injury during this period and that additional reduction of CBF by HV may attenuate the delivery of important energy substrates.^{6,11,29,37,46,47} Yoshida and Marmarou⁵⁸ reported that HV produced relative ischemia in cat brain after fluid percussion injury, showing an increase in brain lactate and inhibition of recovery of the ratio of phosphocreatine to inorganic phosphate. Muizelaar et al³⁹ also demonstrated a loss of brain interstitial bicarbonate buffer after sustained prophylactic HV in rabbits. It has been reported that HV after TBI in animals and humans can reduce CBF to what traditionally have been considered ischemic levels.^{7,23,41} However, defining the ischemic threshold in injured tissue is problematic.^{21,32} Muizelaar, et al³⁷ reported that prolonged HV after TBI in humans may worsen functional outcome raising questions regarding the appropriate indications and timing for the optimum application of HV after TBI. Recent guidelines for the management of severe head injury published in the *Journal of Neurotrauma*⁵ state that "in the absence of intracranial hypertension, HV (PaCO₂ ≤ 35 mm Hg) therapy should be avoided

during the first 24 h after severe TBI....," although, " HV therapy may be necessary for brief periods where there is acute neurologic deterioration..." Consistent with these guidelines, in the setting of acute neurologic deterioration, aggressive HV is used by both emergency and critical care personnel. In addition, in the initial stabilization of the brain injured patient, aggressive HV (planned or iatrogenic) occasionally occurs (in both the pre-hospital and acute care settings). The specific impact of HV during this early low flow period remains to be determined. Despite the availability of well-characterized rodent models of TBI, which reproduce the early posttraumatic reduction in CBF, the effect of aggressive HV on histopathologic and functional outcome have not, to our knowledge, been investigated.

Using a rat model of percussive focal contusion, we hypothesized that aggressive HV, beginning immediately after TBI and continuing for 5 h, would worsen functional outcome, expand the contusion, and promote neuronal death in selectively vulnerable hippocampal neurons.

Materials and Methods

Subjects and Study Groups

All experimental protocols used in this report were approved by the Animal Care and Use Committee of the University of Pittsburgh. Twenty-six virus-free Sprague-Dawley rats (346 ± 5 g) were studied. Food and water were continuously available in their home cages. Rats were randomly assigned after TBI to receive normal ventilation (NV) ($n = 13$, $P_a\text{CO}_2 = 30\text{-}40$ mm Hg) or HV ($n = 13$, $P_a\text{CO}_2 = 15\text{-}25$ mm Hg).

Surgery and Brain Trauma Model

Anesthesia was induced using 4% isoflurane (Anaquest; Memphis, TN) in $\text{N}_2\text{O}:\text{O}_2$ (2:1). The rats were endotracheally intubated and mechanically ventilated. The isoflurane concentration was reduced to 2% followed by sterile surgical placement of a femoral arterial catheter for continuous mean arterial blood pressure (MABP) and arterial blood gas (ABG) monitoring. Penicillin 100,000 U IM (Upjohn, Kalamazoo, MI) and gentamicin 10 mg/kg IM (Elkins-Sinn; Cherry Hill, NJ) were given to minimize infection risk. Pancuronium bromide was administered 0.1 mg/kg/h via the arterial line (Elkins-Sinn; Cherry Hill, NJ) to control ventilation. Core temperature was monitored using a rectal probe.

After stereotactic head positioning (David Kopf; Tujunga, CA), the scalp was retracted, exposing the left parietal bone. A craniotomy was made using a high speed dental drill under a binocular operating microscope. A burr hole was made 5.0 mm anterior and 2.0 mm lateral to the bregma in the left skull and a temperature probe (0.009 in. outer diameter, Physitemp Corp., Clifton, NJ) was inserted through the burr hole 2 mm into the left parietal cortex. The bone flap was left in place and the isoflurane was reduced to 1.0% followed by a 30-min equilibration period. The brain

was maintained at $37 \pm 0.5^\circ\text{C}$. Normal ABGs were achieved in all rats and $P_a\text{O}_2$ was maintained > 70 mm Hg.

TBI was produced using a controlled cortical impact (CCI) device as recently described^{9,24} with minor modifications. Fifteen minutes before CCI, an arterial blood sample was obtained for measurement of ABG, glucose concentration, and hematocrit. The bone flap was removed and a vertical CCI (4 m/s impactor velocity, 2.5 mm deformation depth) was delivered onto the exposed dura overlying the left parietal cortex. The bone flap was replaced and sealed with dental cement and the scalp was resutured.

Study Design

The study protocol was designed to mimic the aggressive use of HV (vs NV) in the immediate post-trauma period in the prehospital as well as early hospital setting. Ten min after CCI, rats were randomized to either the NV group ($n = 13$, $P_a\text{CO}_2$ range, 30-40 mm Hg) or the HV group ($n = 13$, $P_a\text{CO}_2$ range, 15-25 mm Hg). The ventilator was adjusted to maintain normocarbida or hypocarbida for 5 h after CCI. ABGs were obtained at 30 min post-CCI, then hourly. MABP was recorded every 30 min after CCI. Brain and rectal temperatures were recorded every 15 min.

At 5 h after CCI, anesthesia was discontinued. Temperature probes and the femoral artery catheter were removed and the rat was allowed to wean from mechanical ventilation over 1 h and was extubated. Time to extubation was recorded. After extubation, supplemental O_2 was administered for 30 min. Once fully recovered, the rat was returned to its cage with full access to food and water.

Functional Outcome and Behavior Assessment

Beam Balance

Vestibulomotor function was tested using the beam balance test ¹³ in eight rats from each group. One hour before surgery, the rat was placed lengthwise on a 1.5 cm wide beam suspended above ground. The time the rat remained on the beam was recorded (up to 60 s). The rat was then removed from the beam and the procedure was repeated. Rats were considered trained when they remained on the beam for 3 consecutive periods of 60 s. Beam balance tests were also performed daily on d 1-5 after injury. Three trials were recorded and averaged each day for each rat.

MWM

Cognitive function was tested in the same eight rats from each group using a standard variation of the Morris water maze (MWM) paradigm.^{14,34} A pool 180 cm in diameter and 60 cm in depth was painted black and filled with water to a depth of 28 cm. A clear Plexiglas® platform 10 cm in diameter and 26 cm high (2 cm below the water surface) was used as the hidden goal platform. The pool was located in a 2.5 x 2.5 m room with numerous extra-maze cues (e.g., posters, pipes, bookcase) that remained constant throughout the experiment. Testing started 7 d post-CCI in order to avoid motor deficits. Rats underwent four trials per day for 5 consecutive days to assess spatial memory performance. Rats started a trial once from each of the four possible start locations (north, south, east, west). The order of starting location was randomized. The goal platform was positioned 45 cm from the outside wall and was placed in either the northeast, southeast, southwest, or northwest quadrant of the maze. The location of the platform was held constant for each rat. Rats were manually placed in the pool facing the wall and were given a maximum of 120 sec to find the hidden platform. If the rats failed to find the platform within 120 sec, they were placed on the platform by the experimenter. All rats were allowed to remain on the platform for 30 sec before being placed in a heated incubator between trials. There was a 4 min intertrial interval. All data were recorded with a video tracking

system (Poly-Trak Video Tracking System; San Diego Instrument, Inc. San Diego, CA).

Histology

At 14 d after CCI (after completion of all of the functional outcome testing), the rats were anesthetized with 5% isoflurane and killed by perfusion fixation using 10% buffered formalin. The brains were removed and post-fixed at 4°C for a minimum of 1 week and then cryoprotected in sucrose and cut in 10 μ -coronal sections at 1 mm increments from the occipital to the frontal lobe with a cryotome and stained with Cresyl violet.

Contusion Volume

Using a computerized image analysis system (MCID, Imaging Research; St. Catharines, Ontario, Canada) the margin of the contusion was outlined and the sectional area of the contusion at each 1 mm increment was calculated by a blinded observer (MLF). Contusion volume in each rat was calculated as the sum of these sections.

Hippocampal Cell Counting

Neuronal loss in hippocampal regions CA1 and CA3 pyramidal layers was quantified.¹⁰ A coronal section through the dorsal hippocampus underlying the area of contusion, approximately 2.6 mm posterior to bregma was used for analysis for each rat. The regions were visualized at 100x then localized and counted at 400x by a blinded observer (RSBC). Only complete cells with a clearly defined cell body and nucleus were counted. Surviving pyramidal CA1 and CA3 hippocampal neurons were counted in six separate 400x fields for each region in both hemispheres. Sections were excluded if the boundary of the contusion extended into the pyramidal layers of the hippocampus or if fixation artifact precluded accurate counting. Data are

reported as the average number of surviving neurons per high power field for the CA1 and CA3 hippocampal regions in both the ipsilateral and contralateral hemispheres.

Statistical Analysis

Survival was compared between groups using the Fisher's exact test. Between group comparison of physiologic parameters and beam balance and MWM latencies were made using one- or two-way analysis of variance (ANOVA) for repeated measures where appropriate and post-hoc tests with appropriate correction for multiple comparisons. Contusion volume was normally distributed and was compared between groups using the Student's t-test. Hippocampal neuronal survival in CA1 and CA3 was not normally distributed and was compared between groups using the Mann-Whitney Rank Sum test. Significance was defined as $p < 0.05$.

Results

Physiologic Parameters

Baseline and 30-min post-randomization physiologic data are presented for all measured parameters in Table. After randomization, there was a marked increase in pH and decrease in $P_a\text{CO}_2$ in the HV group (vs baseline, $p < 0.05$). HV was also associated with a small (12 mm Hg) increase in $P_a\text{O}_2$ compared with baseline ($p < 0.05$). This difference was attributable to the increased minute ventilation and mean airway pressure in the HV group. At no time were any of the rats hypoxemic ($P_a\text{O}_2 < 100$ mm Hg). The entire time course of arterial $P_a\text{CO}_2$, arterial pH, MABP, and brain temperature after TBI is given for both groups in Figures 1A-D, respectively. Arterial $P_a\text{CO}_2$ and pH differed between groups at all time points after randomization ($p < 0.05$). MABP and brain temperature were similar in both groups.

Five of 26 rats died during the 14-d study. All deaths occurred on the day of injury. Two rats remained unresponsive, were unable to demonstrate any spontaneous respiratory effort for 1 h after discontinuation of anesthesia, and were euthanized. Three rats developed pulmonary edema and/or respiratory distress and died early post-extubation. There were no between group differences in mortality (2/13 vs 3/13, NV vs HV, respectively). There were no between group differences in time to extubation (Table).

Functional Outcome Assessment

Beam balance

There was no difference between groups in motor performance latencies over time ($F_{1,15} = 0.17$, $p < 0.69$, Figure 2). Maximal impairment of performance occurred on d 1 or d 2 in both groups, and eventually returned to baseline. Beam balance performance did not differ significantly between NV and HV groups.

MWM

There was no difference between NV and HV groups in the latency to find the hidden platform in the MWM test ($F_{1,15} = 0.50$, $p < 0.50$, Figure 3). In addition, there was a non-statistically significant trend ($t_{13} = 1.77$, $p < 0.065$) for the HV rats to swim slower than the NV rats (30.8 ± 1.0 vs 35.4 ± 2.1 cm/sec).

Histopathology

Contusion volume

At the injury level selected for this study, contusion was generally restricted to the parietal cortex beneath the impact site. Contusion volume in both groups is shown in Figure 4. There was no difference between groups (27.8 ± 5.1 mm³ vs 27.8 ± 3.3 mm³, NV vs HV) in this outcome parameter.

Hippocampal cell counting

Figure 5 shows the number of surviving neurons/high power field in the CA1 and CA3 regions of the dorsal hippocampus ipsilateral to the contusion. There were no differences in the number of surviving CA1 hippocampal neurons between groups after CCI. There was, however, a further reduction in the number of surviving CA3 neurons in the HV group after CCI compared with the NV group (29.7 [24.2 - 31.7] vs 19.9 [17.0 - 23.7] neurons/high power field, median [25th-75th percentiles], NV vs HV, $p < 0.05$). Neuronal cell counts in the CA1 and CA3 regions in the hemisphere contralateral to the contusion did not differ in either the NV or HV groups (55.3 [52.1 - 59.0] NV CA1 and 57.3 [51.3 - 59.0] HV CA1; 40.0 [36.6 - 41.2] NV CA3 and 38.0 [33.0 - 41.7] HV CA3).

Discussion

In a model of CCI-induced focal contusion in rats, aggressive HV for 5 h after TBI augments neuronal death in CA3 hippocampus ipsilateral to the contusion. However, HV did not worsen motor function or cognitive outcome, as assessed using standard beam balance and MWM paradigms, respectively, and did not increase contusion volume.

Hippocampal CA3 neurons are selectively vulnerable to delayed neuronal death after TBI.^{10,18,48,52,53} The mechanisms underlying this process remain speculative. Potential mechanisms include ischemia, TBI-induced excitotoxicity, apoptosis, and inflammation.^{10,18,48}

Yamakami and McIntosh^{56,57} reported reduced CBF as early as 15 and 30 min after TBI. Using a piglet model of TBI, Pfenninger, et al⁴⁵ reported CBF reduction as early as 5-min post-TBI. Some flows were in the range consistent with ischemia. We have previously demonstrated that the hippocampus and cortex ipsilateral to the impact have marked flow reduction (at least 60%) at 2 h after TBI in the CCI model.²⁴ CBF approaches ischemic levels in the core of the contusion at 2 h after injury. Although we have not evaluated the status of reactivity of the cerebral circulation to changes in PaCO₂ at 2 h after TBI in this model, we have reported that CO₂ reactivity is impaired, although still present (62-71% of baseline) in and around the contusion at 24 h after CCI in rats.¹⁵

HV rapidly reduces CBV and intracranial pressure (ICP).¹¹ In some studies, this intervention has been associated with CBF values consistent with ischemia or brain tissue hypoxia.^{7,11,41,47} After global cerebral ischemia in dogs, HV did not increase neuronal death,⁵⁵ however, brains were assessed at 8 h after reperfusion, and neuronal death may be delayed. While ischemia may be considered a contributing mechanism in the observed augmented neuronal death, ischemia alone is an inadequate explanation for our findings in light of the preservation of CA1 neurons.

Although CA1 neurons are known to be selectively vulnerable to ischemic injury,²² they were not affected by HV in this paradigm. Furthermore, in our model, CA1 neurons are more proximal to the point of impact in the cortex, compared with CA3 neurons. The lack of CA1 neuronal death in light of ischemic and (presumed) anatomic vulnerability mitigates against ischemia and primary injury as putative mechanisms of neuronal death in hippocampus in this model. One limitation in this study is that neuronal counting using traditional histologic methods may underestimate cell loss because of a loss of hippocampal volume.⁵¹ We did not use stereologic methods in this study. However, CA1 neuronal counts did not differ between groups and were equivalent to those observed in shams studied in our laboratory in prior published¹⁰ and unpublished work. In addition, comparisons were only made between injured groups within this study.

HV produces cerebral vasoconstriction and alkalosis.³⁹ Alkalosis exacerbates N-methyl-D-aspartate (NMDA)-receptor mediated neurotoxicity.^{16,17,20,42} As a result of aggressive HV, the rats in our study were quite alkalotic as indicated by arterial pH measurements. Although we did not measure brain pH, a decrease in PaCO_2 immediately reduces brain interstitial pH.³⁹ Although alkalosis appears to have deleterious effects on neurons, acidosis has been shown to have both beneficial and detrimental effects. Giffard et al¹⁷ and others⁵⁴ reported a neuroprotective effect of acidosis via an attenuation of the NMDA receptor activation in vitro. Rosner and Becker⁴⁹ reported a deleterious effect of tissue acidosis after experimental TBI in cats. The spatial distribution of brain pH around the contusion and in the hippocampus has not been determined for either NV or HV conditions in our model.

Finally, the potential effects of HV on other mechanisms such as post-traumatic seizures or axonal injury may contribute to the enhanced vulnerability of CA3 neurons. The lateralization of the deleterious effects also raises the possibility that spreading wave depression may be a component of the neurotoxic milieu after TBI.

in this model of focal contusion.¹⁹ It could also be that the combined effect of alkalosis and further flow reduction by HV is deleterious in regions vulnerable to excitotoxicity such as CA3. Early, aggressive or prophylactic HV, therefore, in the context of reduced CBF, may potentiate excitotoxic mechanisms and augment neuronal death.

Aggressive HV in the early low flow period did not worsen functional outcome or expand the contusion, failing to support a significant portion of our initial hypothesis. Ultimate contusion size, in CCI or other models of focal contusion, is relatively refractory to manipulation by a variety of interventions^{3,10}; however, application of hypothermia particularly prior to injury does reduce contusion volume in CCI.^{12,44} Although we chose rather aggressive HV in an attempt to produce a maximal effect, we did not test the effect of HV on a milder contusion, which may have been more manipulable to secondary insults. The contusion penumbra has not been clearly defined in either of the standard rodent TBI models (CCI or fluid-percussion) for any level of injury. It is possible that selectively vulnerable CA3 hippocampal neurons are the only potential target for a deleterious effect of HV in our model. However, the effect of HV on the survival of neurons in the dentate gyrus or hilus (all vulnerable to TBI)^{9,28} was not assessed.

Hippocampal damage and memory deficits are common after TBI in humans.^{25,27} This study did not reveal any added effect of HV on functional deficits as measured by beam balance and MWM latencies. A number of factors may have contributed to this. Our sample size may have limited statistical power. However, this sample size was adequate to detect the exacerbation of functional deficits by the addition of 30 min of moderate hypoxemia (PaO₂ of 40 mm Hg) in our model.¹⁰ Secondly, the cognitive deficits in this model are modest compared with previous reports.¹⁴ Bilateral hippocampal damage may be necessary to create more marked functional deficits.^{35,36} In addition, CA3 damage may not mediate post-TBI MWM deficits.

Finally, the specific functional outcome paradigm may not have the necessary sensitivity to detect subtle functional deficits. For example, more demanding MWM paradigms have been employed by other investigators.^{26,53} However, in support of the testing strategy used, our hypothesis was that HV would worsen functional deficits.

This study does not completely address the uncommon situation where early after severe head injury marked intracranial hypertension is observed. HV may, in fact, be life saving in the setting of impending herniation. Similarly, we did not measure ICP or titrate ventilation to control cerebral perfusion pressure, and we evaluated only one level of HV and injury severity. We did not attempt to model the clinical scenario of optimal titration of ventilation when ICP is increased. In the clinical setting, some investigators have demonstrated a wide variety of beneficial effects of HV under those conditions, such as homogenization of CBF, normalization of cerebral glucose uptake, and improvement in autoregulation.^{8,40,41} Rather, we chose the worst case scenario, aggressive HV during the early post-trauma period when flow is already low and excitotoxicity is peaking.⁴³ Our study does, however, show that HV is associated with a tangible risk to vulnerable neurons in the CCI model. To our knowledge, this is the first *in vivo* study demonstrating that HV can augment neuronal injury after TBI. This suggests that there is indeed a trade-off associated with this intervention.

Conclusion

We have demonstrated that aggressive, early HV after TBI augments neuronal death in CA3 hippocampus. The further reduction of CBF with HV during the low cerebral blood flow state immediately after severe TBI coupled with alkalosis may increase the vulnerability of selected neurons to traumatic injury. Further studies are needed to delineate the relative contributions of these mechanisms to the

observed effects. The results of this study reinforce the meticulous attention necessary to prevent secondary injury after TBI. A risk in the use of HV is demonstrated.

Acknowledgment

The authors thank Drs. Peter Safar and Walter Obrist for helpful comments. We also thank Raymond Griffith, Erica Baum, and Xiecheng Ma for technical assistance, Linda Amick and Marci Provins for preparing the manuscript, and Francie Siegfried for editorial assistance.

References

1. Artru AA: Reduction of cerebrospinal fluid pressure by hypocapnia: Changes in cerebral blood volume, cerebrospinal fluid volume, and brain tissue water and electrolytes **J Cereb Blood Flow Metab** 7:471-479, 1987
2. Bouma GJ, Muizelaar JP, Stringer WA, et al: Ultra early evaluation of regional cerebral blood flow in severely head injured patients using xenon enhanced computed tomography **J Neurosurg** 77:360-368, 1992
3. Bramlett HM, Dietrich WD, Green EJ, et al: Chronic histopathological consequences of fluid-percussion brain injury in rats: effects of post-traumatic hypothermia **Acta Neuropathol** 93:190-199, 1997
4. Bryan RM, Cherian L, Robertson C: Regional cerebral blood flow after controlled cortical impact injury in rats **J Neurotrauma** 11:104, 1994
5. Bullock RB, Chesnut RM, Clifton G, et al: Guidelines for the management of severe head injury **J Neurotrauma** 13:699-703, 1996
6. Chestnut RM, Marshall LF, Klauber MR, et al: The role of secondary brain injury in determining outcome from severe head injury **J Trauma** 34:216-222, 1993
7. Cold GE: Does acute hyperventilation provoke cerebral oligoemia in comatose patients after acute head injury? **Acta Neurochir** 96:100-106, 1989
8. Cruz J: An additional therapeutic effect of adequate hyperventilation in severe acute brain trauma: Normalization of cerebral glucose uptake **J Neurosurg** 82:379-385, 1995
9. Clark RSB, Kochanek PM, Marion DW, et al: Mild posttraumatic hypothermia reduces mortality after severe controlled cortical impact in rats **J Cereb Blood Flow Metab** 16:253-261, 1996

10. Clark RSB, Kochanek PM, Dixon CE, et al: Early neuropathologic effects of mild or moderate hypoxemia after controlled cortical impact in rats **J Neurotrauma** 14:179-189, 1997
11. Crockard HA, Coppel DL, Morrow WF: Evaluation of hyperventilation in treatment of head injuries **Br Med J** 4:634-640, 1973
12. Deitrich WD, Aloson O, Busto R, et al: Post-traumatic brain hypothermia reduces histopathological damage following concussive brain damage in the rat **Acta Neuropathol** 87:250-258, 1994
13. Dixon CE, Lyeth BG, Povlishock JT, et al: A fluid percussion model of experimental brain injury in the rat **J Neurosurg** 67:110-119, 1987
14. Dixon CE, Ma X, Marion DW: Effects of CDP-Choline treatment on neurobehavioral deficits after TBI and on hippocampal and neocortical acetylcholine release **J Neurotrauma** 14:161-169, 1997
15. Forbes ML, Hendrich KS, Kochanek PM, et al: Assessment of cerebral blood flow and CO₂ reactivity after controlled cortical impact by perfusion magnetic resonance imaging using arterial spin labeling in rats **J Cereb Blood Flow Metab** (in press)
16. Giffard RG, Weiss JH, Choi DW: Extracellular alkalinity exacerbates injury of cultured cortical neurons **Stroke** 23:1817-1821, 1992
17. Giffard RG, Monyer H, Gristine CW, et al: Acidosis reduces NMDA receptor activation, glutamate neurotoxicity and oxygen-glucose deprivation neuronal injury in cortical cultures **Brain Res** 506:339-342, 1990
18. Hicks RR, Smith DH, Lowenstein DH, et al: Mild experimental brain injury in the rat induces cognitive deficits associated with regional neuronal loss in the hippocampus **J Neurotrauma** 10:405-414, 1993

19. Hovda DA, Lee SM, Smith ML, et al: The neurochemical and metabolic cascade following brain injury: Moving from animals to man **J Neurotrauma** 12:903-906, 1995
20. Hurn PD, Koehler RC, Traystman RJ: Extracellular alkalosis reduces recovery from global cerebral ischemia **J Cereb Blood Flow Metab** 15:S201, 1995
21. Jacewicz M, Brint S, Tanabe J, Wang XJ, Pulsinelli WA: Nimodipine pretreatment improves cerebral blood flow and reduces brain edema in conscious rats subjected to focal cerebral ischemia **J Cereb Blood Flow Metab** 10:903-913, 1990
22. Jenkins LW, Moszynski K, Lyeth BG, et al: Increased vulnerability of the mildly traumatized rat brain to cerebral ischemia. The use of controlled secondary ischemia as a research tool to identify common or different mechanisms contributing to mechanical and ischemic brain injury **Brain Research** 477:211-224, 1989
23. Kennealy JA, McLennan, Loudon RG, McLaurin RL: Hyperventilation-induced cerebral hypoxia **Am Rev Resp Dis** 122:407-412, 1980
24. Kochanek PM, Marion DW, Zhang W, et al: Severe controlled cortical impact in rats: Assessment of cerebral edema, blood flow and contusion volume **J Neurotrauma** 12:6 1015-1026, 1995
25. Kotapka MJ, Graham DI, Adams JH, et al: Hippocampal pathology in fatal non-missile human head injury **Acta Neuropath** 83:530-534, 1992
26. Kraemer PJ, Brown RW, Baldwin SA, Scheff SW: Validation of a single-day Morris Water Maze procedure used to assess cognitive deficits associated with brain damage. **Brain Research Bulletin** 39:17-22, 1996
27. Levin HS, Gary HE, Eisenberg HM, et al: Neurobehavioral outcome 1 year after severe head injury **J Neurosurg** 73:699-709, 1990

28. Lowenstein DH, Thomas MJ, et al: Selective vulnerability of dentate hilar neurons following traumatic brain injury: A potential mechanistic link between head trauma and disorders of the hippocampus **J Neurosci** 12:4846-4853, 1992
29. Madsen FF: Changes in regional cerebral blood flow after hyperventilation in the pig with an induced focal cerebral contusion **Acta Neurochir** 106:164-169, 1990
30. Marion DW, Darby J, Yonas H: Acute regional cerebral blood flow changes caused by severe head injuries **J Neurosurg** 74:407-414, 1991
31. Marshall LF, Smith RW, Shapiro HM: The outcome with aggressive treatment in severe head injuries. I. The significance of intracranial pressure monitoring **J Neurosurg** 50:20-25, 1979
32. Meis G, Ishimaru S, Xie , Seo K, Hossman KA: Ischemic Thresholds of Cerebral Protein Synthesis and Energy State Following Middle Cerebral Artery Occlusion in Rat **J Cereb Blood Flow Metab** 11:753-761, 1991
33. Miller JD, Butterworth JF, Gudeman SK, et al: Further experience in the management of severe head injury. **J Neurosurg** 54:289, 1981
34. Morris RGM: Developments of a water maze procedure for studying spatial learning in the rat **J Neurosci Methods** 11:47-60, 1984
35. Morris RGM, Hagan JJ, Rawlins JNP: Allocentric spatial learning by hippocampectomized rats. A further test of the "spatial mapping" and "working memory" theories of hippocampal function **Quart Journ Exp Psychology** 38B:365-395, 1986
36. Moser E, Moser MB, Andersen P: Spatial learning impairment parallels the magnitude of dorsal hippocampal lesions, but is hardly present following ventral lesions **J Neuroscience** 13:3916-3925, 1993

37. Muizelaar JP, Marmarou A, Ward JD, et al: Adverse effects of prolonged hyperventilation in patients with severe head injury: A randomized clinical trial **J Neurosurg** 75:731-739, 1991
38. Muizelaar JP, Marmarou A, DeSelles AA, et al: Cerebral blood flow and metabolism in severely head-injured children. Part 1: Relationship with GCS score, outcome, ICP and PVI **J Neurosurg** 63-71, 1989
39. Muizelaar JP, van der Poel HG, Li ZC: Pial arteriolar vessel diameter and CO₂ reactivity during prolonged hyperventilation the rabbit. **J Neurosurg** 69:923-927, 1988
40. Newell DW, Weber JP, Watson RW, et al: Effect of transient moderate hyperventilation on dynamic cerebral autoregulation after severe head injury **Neurosurg** 39:35-44, 1996
41. Obrist WD, Langfitt TW, Jaggi JL: Cerebral blood flow and metabolism in comatose patients with acute head injury **J Neurosurg** 61:241-253, 1984
42. Ou-Yang YB, Kristian T, Møllergaard P, et al: The influence of pH on glutamate- and depolarization-induced increases of intracellular calcium concentration in cortical neurons in primary culture **Brain Res** 646:65-72, 1994
43. Palmer AM, Marion DW, Botscheller ML, et al: Traumatic brain injury-induced excitotoxicity assessed in controlled cortical impact model **J Neurochem** 61:2015-2024, 1993
44. Palmer AM, Marion DW, Botscheller ML, et al: Therapeutic hypothermia is cytoprotective without attenuating the traumatic brain injury-induced elevations in interstitial concentrations of aspartate and glutamate. **J Neurotrauma** 10: 363-372, 1993
45. Pfenninger EG, Reith A, Breitig D, et al: Early changes of intracranial pressure, perfusion pressure and blood flow after acute head injury. I. An experimental study of the underlying pathophysiology **J Neurosurg** 70:774-780, 1989

46. Raichle ME, Plum F: Hyperventilation and cerebral blood flow **Stroke** 23:566-575, 1972
47. Raichle ME, Posner JB, Posner F: Cerebral blood flow during and after hyperventilation **Arch Neurol** 23:394-403, 1970
48. Rink A, Fung K-M, Trojanowski JQ, Lee VMY, Neugebauer E, McIntosh TK: Evidence of apoptotic cell death after experimental traumatic brain injury in the rat **Amer J Pathol** 147:1575-1583, 1995
49. Rosner MJ, Becker DP: Experimental brain injury: Successful therapy with the weak base, tromethamine **J Neurosurg** 60:961-971, 1984
50. Rosner MJ, Rosner SD, Johnson AH: Cerebral perfusion pressure: management protocol and clinical results **J Neurosurg** 83:949-962, 1995
51. Scheff SW, Baldwin SA: Preparation and staining of tissue for counting neurons. **Kopf Carrier**, David Kopf Instruments, Tujunga, CA, No. 45, pp 1-4, 1996
52. Soares HD, Sinson GP, McIntosh TK: Fetal hippocampal transplants attenuate CA3 pyramidal cell death resulting from fluid percussion brain injury in the rat **J Neurotrauma** 12:1059-1067, 1995
53. Smith DH, Soares HD, Pierce JS, et al: A model of parasagittal controlled cortical impact in the mouse: cognitive and histopathologic effects **J Neurotrauma** 12:169-178, 1995
54. Takadera T, Shumada Y, Mohri T: Extracellular pH modulates N-Methyl-D-aspartate mediated neurotoxicity and calcium accumulation in rat cortical cultures **Brain Res** 572 (1-2) 126-131, 1990
55. Vanicky I, Marsala M, Murar J, et al: Prolonged postischemic hyperventilation reduces acute neuronal damage after 15-minutes of cardiac arrest in the dog. **Neurosci Letters** 135(2): 167-170, 1992

56. Yamakami I, McIntosh TK: Effects of traumatic brain injury on regional cerebral blood flow in rats as measured with radiolabeled microspheres **J Cereb Blood Flow Metab** 9:117, 1989
57. Yamakami I, McIntosh TK: Alterations in regional cerebral blood flow following brain injury in the rat **J Cereb Blood Flow Metab** 11:655-670, 1991
58. Yoshida K, Marmarou A: Effects of tromethamine and hyperventilation on brain injury in the cat **J Neurosurg** 74:87-96, 1991

Table : Physiologic Values ^a

	<i>Normal Ventilation + CCI</i>		<i>Hyperventilation + CCI</i>	
	Baseline	Post-Randomization	Baseline	Post-Randomization
pH	7.39 ± 0.01	7.37 ± 0.01	7.38 ± 0.01	7.53 ± 0.01*
P _a CO ₂ (mm Hg)	36.7 ± 1.1	34.9 ± 0.3	37.2 ± 0.9	20.3 ± 0.7*
P _a O ₂ (mm Hg)	165 ± 6	167 ± 4	168 ± 4	180 ± 3*
Base Deficit (mMol/L)	2.7 ± 3.4	4.2 ± 0.7	-0.6 ± 0.9	4.8 ± 0.6
Serum Glucose (mg%)	189 ± 9	174 ± 6	158 ± 10	152 ± 9
Hct (%)	36.0 ± 2.3	35.0 ± 0.6	32.3 ± 1.5	35 ± 0.6
Time to extubate (min)	N/A	28 ± 6	N/A	29 ± 5
Brain Temperature (°C)	36.7 ± 0.1	37.0 ± 0.0	36.6 ± 0.1	37.0 ± 0.0
Rectal Temperature (°C)	36.5 ± 0.6	37.0 ± 0.0	37.1 ± 0.1	37.1 ± 0.1
Mean Arterial Pressure (mm Hg)	129 ± 4	123 ± 4	129 ± 8	128 ± 3

^a All values expressed as mean ± SEM

* p < 0.05, 30 min post-randomization vs baseline
Controlled cortical impact (CCI)

Figure Legends

Figure 1. Entire time course of (A) PaCO_2 (mm Hg), (B) arterial pH, (C) Mean arterial blood pressure (MABP, mm Hg), and (D) brain temperature ($^{\circ}\text{C}$) in all rats treated with either normal ventilation (NV, \blacktriangle , $n = 13$) or hyperventilation (HV, \blacksquare , $n = 13$) after controlled cortical impact. * $p < 0.05$ for NV vs HV. Data are mean \pm SEM.

Figure 2. Mean beam balance performance latencies (mean \pm SEM, in sec) in rats before and on d 1-5 after controlled cortical impact (CCI: 4 m/s, 2.5 mm cortical deformation depth). Analysis of variance with repeated measures revealed no difference in duration of balance maintained between the two groups. (\blacktriangle , normal ventilation, $n = 8$; \blacksquare , hyperventilation, $n = 8$). Data are mean \pm SEM.

Figure 3. Morris water maze performance latency to find a hidden platform (mean \pm SEM, in sec) by rats on d 7-11 after controlled cortical impact. There was no between group difference (\blacktriangle , normal ventilation, $n = 8$; \blacksquare , hyperventilation, $n = 8$) when performances were compared using ANOVA with repeated measures. Data are mean \pm SEM.

Figure 4. Graph depicting mean lesion area (left y-axis, mm^2) vs distance from occiput (mm) measured 14 d after controlled cortical impact (normal ventilation NV, open bars, $n = 11$ / hyperventilation, HV, closed bars, $n = 10$). Contusion volume (mm^3) was calculated as the sum of these areas in each group and is depicted as cumulative volume (right y-axis) in the NV, \blacktriangle , and HV, \blacksquare , groups. There was no difference between groups in contusion volume (27.8 ± 5.1 vs $27.8 \pm 3.1 \text{ mm}^3$ NV vs HV, mean \pm SEM).

Figure 5. Box plots representing the number of surviving CA1 and CA3 hippocampal neurons in coronal brain sections through the center of the lesion in the hemisphere ipsilateral to the contusion. Cells were counted 14 d after injury. The median line is placed within the shaded 25th - 75th range. There was a reduction in the number of surviving CA3 hippocampal neurons after injury comparing normal ventilation (NV) and hyperventilation (HV) groups (29.7 [24.2 - 31.7] vs 19.9 [17.0 - 23.7], cells/high power field (hpf), $*p < 0.05$, Mann-Whitney rank sum test).

Figure 1
Forbes et al




Figure 2
Forbes et al




Figure 3
Forbes et al




Figure 4
Forbes et al




Figure 5
Forbes et al

